

# Having a Colonoscopy with Klean Prep

## Patient Information Leaflet Gastroenterology Department



### Options available

If you'd like a large print, audio, Braille or a translated version of this leaflet then please call:

**01253 955588**

Our Four Values:

People Centred

Positive

Compassion

Excellence

## **What to expect on the day**

A Colonoscopy is a test, which allows a Doctor to look directly at the whole of the large bowel (Colon). In order to do this a flexible tube the thickness of a finger with a light on the end of it is carefully passed through the anus into the rectum and large bowel. This can be uncomfortable as bends in the colon are negotiated. The Colonoscope is connected to a television system where pictures can be seen. It is important you have a clean bowel in order for us to complete this test.

## **Benefits**

By examining your bowel the cause of your symptoms may be found, sometimes we are able to treat them there and then. Pictures and samples (biopsies) may be taken and sometimes polyps (small growths in the bowel) can be removed or other treatments performed.

## **Risks**

A Colonoscopy is a safe examination but rarely a complication such as bleeding or perforation can occur (1 in 1500) or if polyps removed (1 in 200).

## **Alternative Treatments**

The alternative to a Colonoscopy is a barium enema, a type of x-ray, which does not include treatment and cannot provide biopsies therefore you may still need to have a Colonoscopy at a later date.

## **Please Note**

Your appointment time takes into account the time required to admit you to the unit therefore you should not expect to immediately go through for the test when called. You should expect to be with us for between 2-3 hours (including recovery time).

## What to expect on the day

- Before the procedure a nurse will take you to a room and explain the test and ask you some questions. Some of this may be done by phone prior to the day of examination.
- You will then be asked to sign a consent form and change into a hospital gown.
- A small cannula will be placed in your arm to enable us to give you medication for the test, you will then be taken to a trolley and into the room for your test after a short wait.
- In the endoscopy room a sedative is given through the cannula. This is not a general anaesthetic, it will relax you but you will be aware of what is going on. This may cause loss of memory and judgement for some time.
- Following the test you will be moved to the recovery area for about one hour to rest.
- You may feel bloated with wind like discomfort. This is normal and usually settles quickly. If you have a biopsy taken or a polyp removed it is normal to pass small spots of blood from your bottom for a day or two.
- We will ask you not to drive, drink alcohol, operate machinery or sign legal documents for 24 hours after the tests.
- You will need someone to accompany you home because of the effects of the sedation and also someone at home with you for 12 hours following the procedure.
- You will be given the results of the test before leaving the unit (except biopsies).

**N.B. Please leave all valubles and jewellery at home where possible as we can not look after these for you.**

## **Klean Prep Instructions**

Klean-Prep works by giving you diarrhoea. This is unpleasant but necessary and may last up until your appointment. Easy access to a toilet will be required for the most of the day.

**Note: If you have a Colostomy or Ileostomy do not take the bowel preparation - contact the telephone number on top of your appointment letter for advice.**

### **Seven days before the Examination**

Stop taking iron tablets

If you have chronic constipation (stools infrequent, hard and difficult to pass) take 2 Senna 7.5mg tablets at night for 1 week before beginning your bowel preparation.

### **Two days before the Examination**

Stop taking any medicines that might constipate you for example codeine, co-phenotrope (Lomotil®), loperamide (Imodium®) or Kaolin and Morphine mixture.

Stop taking bran and high roughage foods.

It is important that you follow a low residue diet as part of your bowel preparation the day before taking Klean-Prep.

You will find details of low residue foods on page 8.

## **If your appointment is in the morning**

### **On the Day Before the Examination**

**1pm** - Have a light lunch, avoiding all foods containing roughage, including wholemeal bread and vegetables. After this keep to a liquid diet: Clear soup, Bovril, Oxo, fizzy drinks, clear fruit juices, black tea or coffee (no red liquids).

**3pm** - Empty the contents of sachet one of Klean-Prep into 1 litre (1  $\frac{3}{4}$  pints) of water. Stir until the solution is clear. Drink 1 glassful ( $\frac{1}{4}$  litre or  $\frac{1}{2}$  pint) of Klean-Prep about every 15 minutes until you have drunk it all.

**4pm** - Take sachet two (mixed as above)

**5pm** - Rest period of one hour

**6pm** - Take sachet three (mixed as above)

**7pm** - Take sachet four (mixed as above)

This is an example of timings to take your Klean Prep please refer to manufacturers leaflet for further instructions.

The solutions from all 4 sachets should be drunk within 4 to 6 hours.

**If you have not had a bowel movement after taking two sachets, wait until you go to the toilet before you continue taking the Klean- Prep.**

**If you have still not had a bowel movement by the next morning please phone for further advice.**

## **If your appointment is in the afternoon**

### **On the Day Before the Examination**

**4pm** - Have a light meal, avoiding all foods containing roughage, including wholemeal bread and vegetables. After this keep to a liquid diet: Clear soup, Bovril, Oxo, fizzy drinks, clear fruit juices, black tea or coffee (no red liquids).

**6pm** - Empty the contents of sachet one of Klean-Prep into 1 litre (1  $\frac{3}{4}$  pints) of water. Stir until the solution is clear. Drink 1 glassful ( $\frac{1}{4}$  litre or  $\frac{1}{2}$  pint) of Klean-Prep about every 15 minutes until you have drunk it all.

**7pm** - Take sachet two (mixed as above)

Rest overnight

### **On the Day of Examination**

**8am** - Take sachet three (mixed as above)

**9am** - Take sachet four (mixed as above)

This is an example of timings to take your Klean-Prep please refer to manufacturers leaflet for further instructions.

The solutions from all 4 sachets should be drunk within 4 to 6 hours.

**If you have not had a bowel movement after taking 2 sachets, wait until you go to the toilet before you continue taking Klean-Prep.**

**If you have still not had a bowel movement by the next morning please phone for further advice.**

## **Important notes on Klean Prep**

### **Possible Side effects of Klean-Prep**

Nausea, vomiting, bloating and stomach cramps are common side effects. If you experience any of these try taking Klean-Prep more slowly.

Rarely people are allergic to Klean Prep and may get an itchy skin rash (nettle rash or hives) If any other symptoms occur please phone the number on top of your appointment letter, if out of hours seek medical advice.

### **Female Patients**

If you are taking the oral contraceptive pill, diarrhoea can make it less effective. Continue taking the pill but use other precautions for the rest of that cycle.

### **Important:**

**No alcohol should be drunk on the day before and the day of the examination.**

**If you have not had a bowel movement prior to taking Klean-Prep or if you have any abdominal swelling or discomfort please DO NOT take Klean-Prep without first calling the Gastroenterology Department for further advice.**

## **What to eat for your Low Residue Diet**

**Cereals:** Crisped rice cereal, Cornflakes (no bran).

**Fats:** Butter, margarine (use sparingly).

**Eggs:** Boiled, poached.

**Meat / Fish:** Minced or well cooked, tender lean Beef, Lamb, Ham, Veal, Pork, Poultry, Fish, Shellfish.

**Rice:** Plain boiled white rice.

**Pasta:** Macaroni, spaghetti, Noodles.

**Bread:** White bread / toast.

**Potatoes:** Boiled, creamed, mashed or plain, baked (no skins).

**Cheese:** Cream, cottage or cheese sauce.

**Dessert:** Clear jelly (not red Jelly).



## **Instructions for Diabetics**

The necessary change in your diet may affect your diabetes and your blood glucose (sugar). However, do not worry as this will not do any long-term harm. On the day before your examination you may be at the risk of having a "hypo" (too low blood sugar). When on clear fluid diet you can have sweet drinks (clear fluids only such as Lucozade) every 2-3 hours during the day to avoid the problem.

We aim not to admit patients during the 2-day preparation period before their examination unless their diabetes is particularly unstable, and we hope the enclosed directions are helpful.

If you have any queries then contact the Diabetic Liaison Nurse. You should consider seeking advice if your diabetic control becomes unsatisfactory.

## **Patients on Hypoglycaemic Tablets**

### **On the Day Before the Examination.**

If applicable you should carry out your usual pre-meal blood sugar monitoring

- Continue on your usual tablet regime.
- Monitor your blood sugar, every 4-6 hours.
- If your blood sugar falls to around 5mmol/l or less, or if you experience hypoglycaemic symptoms, take 10-20g of carbohydrate in sugary drinks.

- If your blood sugars are too high at mealtimes (say 12mmol/l or more), despite your usual doses of tablets, you may need to reduce your quantity of sugary drinks.

### **On the Day Of the Examination:**

Do not take your tablets on the morning of the examination, but you can take them with some food after the examination.

### **Patients on Insulin on the Day Before the Examination.**

#### **Twice daily insulin regime:**

If taking a twice-daily insulin regime (before breakfast and before evening meal) (e.g. Insulatard insulin, Novomix 30, Humulin I, Humalog Mix 25, Humalog Mix 50, Humulin M3) reduce both doses by 20-25% (but no more).

#### **Four times per day Insulin regime:**

- If taking a 4 times a day Insulin regime, then reduce the meal times doses of quick acting soluble insulin (Insulin (Actrapid, Humalog, Novorapid, Humulin S, Apidra (but no more).
- The bedtime dose of long acting insulin (Lantus, Levemir, Abasaglar, Toujeo and Tresiba) remains unchanged and should be taken on the evening before the procedure.

#### **All patients on Insulin:**

- Blood Glucose testing should be done 2-4 hourly, in order to prevent the development of hypoglycaemia.
- You can take 10-20g of carbohydrate in sugary drinks if your blood sugar falls to around 5 mmol/L or less, or if experiencing hypoglycaemia.

You may need to be flexible with the amount of sugary drinks ingested; for instance, take smaller quantities than suggested above if the blood sugars are too high at mealtimes (say 12 mmol/L or more), despite your usual doses of insulin.

### **On the Day Of the Examination**

If your examination is in the morning: do not take your usual morning dose of insulin.

Bring your insulin with you so that you can take it after the procedure, together with some food (after checking your blood sugar).

If your examination is in the afternoon: take a reduced dose of insulin (reduce doses by 25-50%) at breakfast time, together with a glucose drink. Bring your insulin with you so that you can take it after the procedure, together with some food (after checking your blood sugar).

### **Sugary drinks include:**

- Glucose drinks - 55mls contains 10g of carbohydrate
  - Fizzy lemonade or coke - 100mls contains 10g of carbohydrate
- 1 glass (180-200mls) of glucose drinks or 2 glasses of lemonade/coke can be sipped slowly at meal times to replace your usual dietary carbohydrate.

Remember, if you have queries then contact the Diabetic Liaison Nurses for advice.

**Please bring a list of your medications with you on the day of test.**

**References:** There are too many references to add to the leaflet and therefore a full list of references can be obtained from the policy coordinator. Telephone: 01253 953397

## Useful contact details

The Gastroenterology Unit entrance is located via the Woodlands Car Park off East Park Drive (near the golf course). There is also access via an external door off the main corridor. If you have any problems before or after your test you can contact the Gastroenterology Unit on 01253 953043. If out of hours contact your GP or your local A&E.

Useful websites:  
Patient UK Colonoscopy

Hospital Switchboard: **01253 300000**

## Patient Relations Department

The Patient Relations Department offer impartial advice and deal with any concerns or complaints the Trust receives. You can contact them via:

Tel: **01253 955589**  
email: [patient.relations@bfwh.nhs.uk](mailto:patient.relations@bfwh.nhs.uk)

You can also write to us at:

**Patient Relations Department, Blackpool Victoria Hospital,  
Whinney Heys Road, Blackpool FY3 8NR**

Further information is available on our website: [www.bfwh.nhs.uk](http://www.bfwh.nhs.uk)

## References

This leaflet is evidence based wherever the appropriate evidence is available, and represents an accumulation of expert opinion and professional interpretation.

Details of the references used in writing this leaflet are available on request from: **Procedural Document and Leaflet Coordinator 01253 953397**

**Approved by:** Medicine Management  
**Date of Publication:** 20/04/2017  
**Reference No:** Ic00010667 - PL/540 (v2)  
**Author:** Claire Lester  
**Review Date:** 01/04/2020